

COVID-19 Vaccines

Why get the COVID-19 vaccine?

The vaccines have safely protected millions of people from getting very sick, hospitalized, or dying from COVID-19. The protection you gain from the vaccine is less risky than getting COVID.

COVID-19 vaccines and booster shots for most people.

The Pfizer vaccine is available for both children and adults:

- Children ages 6 months to 4 years can get 2 doses, 3 weeks apart, and a 3rd dose 8 weeks later. All 3 doses are required to complete the series.
- Everyone 5 and older can get 2 doses, 3 weeks apart, and a booster dose 5 months later.

The Moderna vaccine is available for both children and adults:

- Children ages 6 months - 17 years can get 2 doses, 4 weeks apart.
- Adults 18 and older can get 2 doses, 4 weeks apart, and a booster dose 5 months later.

The Johnson and Johnson (J&J) vaccine is available for adults:

- Adults 18 and older can get 1 dose and then a booster shot after 2 months. People who got J&J as their primary and booster dose, can get a second booster shot 4 months later.

If you are immunocompromised, you should get extra doses.

People who should get an extra doses are people who:

- Have a weakened immune system
- Have had an organ or stem cell transplant
- Have cancer
- Have HIV.
- Have chronic conditions such as diabetes or heart, liver, kidney, and lung diseases.

If you think you might be in this group, please talk with a doctor or medical provider. People in this group can get extra primary and booster doses. Vaccine schedules for people who are immunocompromised can be found on the next page.

If you are aged 50 years or older, you should get a second booster.

Adults ages 50 and older, both immunocompromised and not immunocompromised, can choose to get a second booster shot 4 months after their first booster shot.


Mixing and matching booster shots.

Adults 18 and older can get any brand of vaccine as their first booster shot. For the best protection, it is recommended to get either Pfizer or Moderna. Children cannot mix and match vaccines and not all children can get a booster at this time.


See a schedule for each vaccine on the next page.

Vaccine Schedule for Most People

Pfizer
(Ages 6 months to 4 years)



Pfizer
(Ages 5 and older)



4 months

2nd Booster




People ages 50 and older should get a second booster.

Moderna
(Ages 6 months to 17 years)



Moderna
(Ages 18 and older)



4 months

2nd Booster



People ages 50 and older should get a second booster.

Johnson & Johnson
(Ages 18 and older)



4 months

2nd Booster



- People ages 50 and older should get a second booster.
- People ages 18 and older who got J&J as their primary and booster dose, can get a second booster that is either Pfizer or Moderna.

Vaccine Schedule for Immunocompromised

Pfizer
(Ages 6 months to 4 years)



Pfizer
(Ages 5 to 11)



Pfizer
(Ages 12 and older)



Moderna
(Ages 6 months to 17 years)



Moderna
(Ages 18 and older)



Johnson & Johnson
(Ages 18 and older)

