# NEW MEXICO COVID-19 CASES UPDATE STATEWIDE AND COUNTY-LEVEL TRENDS


### December 19, 2022


Unless stated otherwise, all data reported here exclude cases who are from out-of-state and cases who are detainees in Federal Immigration and Customs Enforcement (ICE) facilities.

Total Cases	Cases in the Last 7 Days
655,720 <sup>1</sup>	2346

## SECTION 1: STATEWIDE AND COUNTY-LEVEL CASES


New Mexico cases by date of specimen collection with 7 day moving average







24796-30386

35681-45840








Note: Data updated 12/20/2022 and downloaded from https://coronavirus.jhu.edu/. For U.S. interstate comparisons, the methodology used here is slightly different than methodologies used in other NMDOH COVID-19 reports.

#### Percentage of all emergency department (ED) visits that were Coronavirus-like illness (CLI) and Influenza-like illness (ILI) related


These visualizations are populated from data in New Mexico's Syndromic Surveillance Database. Initial patient encounter information is usually received within 24 hours, but clinical documentation is continuously being updated as it is identified throughout the patient encounter and hospital coding process.



- ILI CC DD

**CLI CC with CLI DD and Coronavirus DD** includes ED encounters with chief complaint consisting of fever and cough, shortness of breath, or difficulty breathing, while also including COVID-19 associated discharge diagnoses codes. The CLI definition excludes known influenza related ED visits coded with related influenza discharge diagnosis.

**ILI CCDD** includes ED encounters with chief complaint consisting of fever and cough, while also including ILI and influenza related discharge diagnoses.




### Cumulative number of cases and recovered cases by county


County	Cumulative Cases	Cumulative Recovered
Bernalillo	192881	186076
Catron	518	478
Chaves	24222	23576
Cibola	9044	8666
Colfax	2756	2661
Curry	14888	14450
De Baca	728	706
Dona Ana	74916	72344
Eddy	18949	18420
Grant	9570	9031
Guadalupe	1327	1270
Harding	101	97
Hidalgo	1295	1260
Lea	22238	21396
Lincoln	6413	6204
Los Alamos	4520	4395
Luna	7866	7608
McKinley	32202	30828
Mora	976	938
Otero	16600	15916
Quay	2478	2360
Rio Arriba	13034	12682
Roosevelt	5996	5788
San Juan	51383	49130
San Miguel	7506	7265
Sandoval	43953	42499
Santa Fe	39444	38287
Sierra	2452	2312
Socorro	4630	4425
Taos	7177	6941
Torrance	3273	3116
Union	924	901


# **SECTION 2: TESTING**

#### Testing rate by U.S. States



Note: Data downloaded 12/20/2022 and downloaded from https://beta.healthdata.gov/National/COVID-19-Community-Profile-Report/gqxm-d9w9. For U.S. interstate comparisons, the methodology used here is slightly different than methodologies used in other NMDOH COVID-19 reports.





Note: Data downloaded 12/20/2022 and downloaded from https://beta.healthdata.gov/National/COVID-19-Community-Profile-Report/gqxm-d9w9. For U.S. interstate comparisons, the methodology used here is slightly different than methodologies used in other NMDOH COVID-19 reports. States colored gray in the map are missing data this week.

- COVID-19 data:
- New Mexico Electronic Disease Surveillance System (NM-EDSS), Infectious Disease Epidemiology Bureau, Epidemiology and Response Division, New Mexico Department of Health.
- Salesforce/MTX COVID-19 Case Investigation Platform.
- Population Estimates: University of New Mexico, Geospatial and Population Studies (GPS) Program.
- Age-adjustment: US 2000 Standard Population Weights

#### Data Notes

- The data reported in this weekly update may not match the daily numbers that are reported in the New Mexico Department of Health (NMDOH) press releases and/or the NMDOH COVID-19 data dashboard. This may be due to variation in the date and time of data extraction from NM-EDSS, corrections after quality assurance review, and differences in the exclusion criteria.
- New Mexico Electronic Disease Surveillance System (NM-EDSS). Disease incidence data are derived from reports of notifiable infectious diseases. NMDOH relies on health care providers, laboratories, hospitals, clinics, institutions and individuals to report suspected and confirmed notifiable infectious diseases in accordance with New Mexico Administrative Code 7.4.3.13. Under-reporting can occur due to of lack of awareness about reporting requirements or lack of compliance with those requirements. Not all cases of infectious diseases can be detected for various reasons including lack of access to health care services, lack of laboratory testing or concerns about confidentiality. Specific and standardized national case definitions are used to classify disease reports by case status.
- New Mexico Population Estimates. All population estimates apply to July 1 of 2019. Estimates include decimal fractions. The sum of population subgroup estimates may not exactly equal the overall state population estimate due to rounding error. Population estimates for previous years are occasionally revised as new information becomes available. When publishing trend data, always be sure that your rates for earlier years match current rates on NM-IBIS that have been calculated with the most up-to-date population estimates.
- **Race/Ethnicity.** Race/Ethnicity are reported as a single variable according to the selection of the case. Any case who is Hispanic is in the Hispanic category and all other races are non-Hispanic.
- **Gender** refers to a person's internal sense of being male, female, some combination of male and female, or neither male nor female. Sex refers to the biological anatomy of an individual's reproductive system, and secondary sex characteristics.
- Case rate per 100,000 population. A basic measure of disease-specific case frequency is a rate, which takes into account the number of cases and the population size. It is helpful in making public health decisions for a given population, relative to another population regardless of size.
- Age-adjusted case rate per 100,000 population. The age-distribution of a population (the number of people in particular age categories) can change over time and can be different in different geographic areas. The use of age-adjusted rates permits a valid comparison among populations. It ensures that the differences in cases from one population to another are not due to differences in the age distribution of the populations being compared.